

Divergence of CAT(0) Cube Complexes and Coxeter Groups

Ivan Levcovitz

Abstract

We provide geometric conditions on a pair of hyperplanes of a CAT(0) cube complex that imply divergence bounds for the cube complex. As an application, we classify all right-angled Coxeter groups with quadratic divergence and show right-angled Coxeter groups cannot exhibit a divergence function between quadratic and cubic. This generalizes a theorem of Dani-Thomas that addressed the class of 2-dimensional right-angled Coxeter groups. As another application, we provide an inductive graph theoretic criteria on a right-angled Coxeter group's defining graph which allows us to recognize arbitrary integer degree polynomial divergence for many infinite classes of right-angled Coxeter groups. We also provide similar divergence results for some classes of Coxeter groups which are not right-angled.

1 Introduction

In this article we study the divergence of CAT(0) cube complexes and apply our results to the class of right-angled Coxeter groups. We also provide results regarding the divergence of Coxeter groups which are not right-angled.

Given a metric space X and a positive number r , the divergence function $Div(X, r) = Div(X)$ is the supremum over all lengths of minimal paths, which avoid a ball of radius r , connecting two points that are distance roughly r apart. One may roughly think of the divergence function as a measure of the best upper bound on the rate a pair of geodesic rays can stray apart from one another. For a finitely generated group G , $Div(G)$ is the divergence function applied to the Cayley graph of G endowed with the word metric.

Groups which are δ -hyperbolic all exhibit at least exponential divergence. On the other end of the spectrum, \mathbb{Z}^n displays linear divergence for $n \geq 2$. Gromov conjectured that nonpositively spaces should exhibit either linear or at least exponential divergence [Gro93]. To the contrary, many important classes of groups, several of which are CAT(0), contain groups of quadratic divergence. For instance, the class of 3-manifold groups, the mapping class group of a closed surface of genus $g \geq 2$, right-angled Artin groups and right-angled Coxeter groups have been shown to contain groups of quadratic divergence [Ger94a] [KL98] [Beh06] [BC12] [DT15].

More recently, there have been some examples of CAT(0) groups exhibiting polynomial divergence of any positive integer degree [BD14] [BH16] [DT15] [Mac13]. Additionally, there are even constructions of more exotic infinitely presented groups with divergence function not a polynomial [GS] [OOS09].

Right-angled Artin groups have played a central role in contemporary mathematics (see for example [Wis11] [Ago13]). In terms of their divergence, these groups satisfy a certain trichotomy: each right-angled Artin group either exhibits linear, quadratic or infinite divergence with these occurrences classified by simple properties of the group’s defining graph [BC12]. 3-manifolds exhibit a similar trichotomy as well [Ger94b].

Right-angled Coxeter groups form an important class of groups which act geometrically on CAT(0) cube complexes. Associated to any simplicial graph Γ is a right-angled Coxeter group, W_Γ , whose presentation consists of an order 2 generator for each vertex of Γ with the relation that two generators commute if there is an edge between the corresponding vertices of Γ . It is true that every right-angled Artin group is finite index in some right-angled Coxeter group [DJ00], therefore as divergence is a quasi-isometry invariant, the class of right-angled Coxeter groups contains groups of linear, quadratic and infinite divergence as well. However, even more is true for these groups.

In fact, for any positive integer degree, Dani-Thomas surprisingly provide an example of a 2-dimensional right-angled Coxeter group exhibiting polynomial divergence of the given degree [DT15]. This raises the question of which divergence functions are possible for right-angled Coxeter groups. Additionally, there is the question of which properties of right-angled Coxeter groups give rise to their broader spectrum of divergence functions and how can these properties be recognized through these groups’ defining graphs.

The class of groups that act geometrically on a CAT(0) cube complex is vast and includes both right-angled Artin groups and right-angled Coxeter groups. More generally, we ask which properties of CAT(0) cube complexes give rise to different divergence functions.

The Rank Rigidity Theorem shows the existence of a rank one isometry in an irreducible, essential, locally compact CAT(0) cube complex with cocompact automorphism group [CS11] (the result actually holds under more general assumptions as well). As a consequence, the divergence of these spaces is either linear or at least quadratic [Hag13b].

We prove for the case of right-angled Coxeter groups, there is an additional gap between quadratic and cubic divergence, and we classify exactly which right-angled Coxeter groups exhibit quadratic divergence.

Theorem 7.4 *Suppose the graph Γ is not a non-trivial join. The right-angled Coxeter group W_Γ exhibits quadratic divergence if and only if Γ is CFS. If Γ is not CFS, then the divergence of W_Γ is at least cubic.*

The CFS condition (“constructed from squares”) is a purely graph-theoretic condition which can be computationally checked. We note that the divergence of W_Γ is linear if and only if Γ is a non-trivial join [BFRHS]. For the case when Γ does not contain triangles, the above theorem is a result of Dani-Thomas [DT15]. Such groups are precisely those whose Davis complex, a natural CAT(0) space a Coxeter group acts on, is 2-dimensional. Theorem 7.4 thus generalizes Dani-Thomas’s result to right-angled Coxeter groups of arbitrary dimension.

An important application of Theorem 7.4 is to the theory of random right-angled Coxeter groups. Let $\Gamma(n, p(n))$ be a random n -vertex graph containing an edge between a given pair of vertices with probability $p(n)$. A random right-angled Coxeter group is simply the right-angled Coxeter group defined by a random graph. Behrstock–Falgas-Ravry–Hagen–Susse [BFRHS] give a threshold theorem for when a random graph is

CFS with probability 1. Combining their result with Theorem 7.4, we obtain a threshold function for the transition between quadratic to at least cubic divergence in random right-angled Coxeter groups.

Theorem 1.1 (Behrstock–Falgas–Ravry–Hagen–Susse, Levcovitz). *Suppose $p(n)$ is a probability density function bounded away from 1 and let $\epsilon > 0$. Let $\Gamma = \Gamma(p(n), n)$ be a random graph. If $p(n) > n^{-\frac{1}{2}+\epsilon}$, then the right-angled Coxeter group W_Γ asymptotically almost surely exhibits quadratic divergence. If $p(n) < n^{-\frac{1}{2}-\epsilon}$, then W_Γ asymptotically almost surely exhibits at least cubic divergence.*

We note that a random right-angled Coxeter group given as above asymptotically almost surely has dimension larger than two. Thus the generality of Theorem 7.4 is needed to obtain Theorem 1.1.

Strongly thick metric spaces of order d form an important class of spaces which can be constructed through a d -step inductive gluing procedure, with initial pieces of linear divergence. An important consequence is that these spaces must have divergence bound above by a polynomial of degree $d + 1$ [BD14]. There are not many general results in the opposite direction giving lower bounds on divergence, and a goal of this article is to introduce criteria which imply such lower bounds. Having such criteria then allows us to give the exact divergence, up to an equivalence of functions, for many spaces.

We apply the following strategy to study the divergence in CAT(0) cube complexes. First, we define the *hyperplane divergence function* that, for each pair of non-intersecting hyperplanes, gives the length of a shortest path between these hyperplanes that avoids a ball of radius r about a basepoint. We then give conditions on a pair of non-intersecting hyperplanes that imply a lower bound on their corresponding hyperplane divergence function. The proof for these lower bounds involve the use of disk diagrams. Finally, we show how the hyperplane divergence function for a pair of such hyperplanes actually implies a stronger lower bound on the divergence of the entire CAT(0) cube complex:

Theorem 5.10 *Let X be an essential, locally compact CAT(0) cube complex with cocompact automorphism group. Suppose $H\text{Div}(\mathcal{Y}, \mathcal{Z}) \succeq F(r)$ for a pair of non-intersecting hyperplanes \mathcal{Y} and \mathcal{Z} in X . It then follows that $\text{Div}(X) \succeq rF(r)$.*

Consequently, this process reduces the problem of finding a lower bound on divergence to finding a pair of hyperplane with certain separation properties. Through this strategy, we prove the following theorem which gives lower bounds on divergence as a consequence of the existence of certain types of pairs of non-intersecting hyperplanes.

Theorem 5.3 and 6.2 *Suppose X is an essential, locally compact CAT(0) cube complex with cocompact automorphism group. Let \mathcal{Y} and \mathcal{Z} be non-intersecting hyperplanes in X .*

1. *If \mathcal{Y} and \mathcal{Z} are k -separated, then $\text{Div}(X)$ is bounded below by a quadratic function.*
2. *If \mathcal{Y} and \mathcal{Z} are k -chain separated, then $\text{Div}(X) \succeq \frac{1}{2} \log_2(\log_2(R))R^2$.*
3. *If X contains a pair of degree d k -separated hyperplanes, then $\text{Div}(X)$ is bound below by a polynomial of degree $d + 1$.*
4. *Suppose X has k -alternating geodesics. If \mathcal{Y} and \mathcal{Z} are symbolically k -chain separated then $\text{Div}(X)$ is bounded below by a cubic function.*

The aforementioned classification of quadratic divergence in right-angled Coxeter groups is an application of 4 above. Furthermore, as an application of 3 we give graph-theoretic criteria which imply polynomial lower bounds on divergence of right angled Coxeter groups.

Theorem 7.9 *Suppose the graph Γ contains a rank n pair (s, t) , then $\text{Div}(W_\Gamma)$ is bound below by a polynomial of degree $n + 1$.*

Here a rank n pair (s, t) is a pair of non-adjacent vertices $s, t \in \Gamma$ which satisfy a certain inductive graph-theoretic criteria. By the above theorem and the machinery of thickness, we provide exact bounds on the divergence of a wide range of right-angled Coxeter groups. The above theorem, in particular, applies to the examples given in [DT15].

Finally, we explore the divergence in the setting of Coxeter groups (not necessarily right-angled). Given an edge-labeled simplicial graph Γ there is a corresponding Coxeter group W_Γ . An adaptation of our techniques allow us to prove results in this general case. For instance, we provide the following polynomial lower bound.

Theorem 8.4 *Let Γ be an even triangle-free Coxeter graph. Suppose (u, v) is a rank n pair, then the divergence of the Coxeter group W_Γ is bound below by a polynomial of degree $n + 1$ in r .*

By the thickness machinery and the above theorem, for any positive integer degree, we can conclude there are infinite classes of Coxeter groups which are not right-angled and which have polynomial divergence of that degree. This shows the existence of higher degree polynomial divergence in the general class of Coxeter groups is abundant.

Theorem 8.4 is actually proven in a more general setting as we only need Γ to be triangle-free and even for some neighborhood of the vertex u . For a precise statement see Theorem 8.4 of Section 8. For an edge-labeled graph Γ representing a Coxeter group, we let $\hat{\Gamma}$ denote the graph obtained by homotoping odd labeled edges of Γ to a point. We prove the following:

Theorem 8.7 *Let Γ be a Coxeter graph. If the diameter of $\hat{\Gamma}$ is larger than 2, then W_Γ has at least quadratic divergence.*

In particular, the above theorem shows that if W_Γ is an even Coxeter group where Γ has diameter larger than 2, then the divergence of W_Γ is at least quadratic.

The paper is organized as follows. Section 2 provides general background material, including a background on divergence, $\text{CAT}(0)$ Cube Complexes and Coxeter groups. Section 3 provides necessary background on disk diagrams in $\text{CAT}(0)$ cube complexes. In this section we set the notation and results regarding disk diagram structures that are used throughout the article.

In section 4, we introduce several notions of separation for a pair of non-intersecting hyperplanes in a $\text{CAT}(0)$ cube complex. The consequences of these separation properties on the divergence of a $\text{CAT}(0)$ cube complex are explored in section 5 and 6. The hyperplane divergence function is defined there as well.

In section 7 we apply the results obtained for $\text{CAT}(0)$ cube complexes to the setting of right-angled Coxeter groups. Finally, in section 8, we explore the divergence of Coxeter groups which are not necessarily right-angled.

Acknowledgements: I would like to especially thank my advisor Jason Behrstock for countless discussions and guidance in writing this paper. I would also like to thank

Mark Hagen for many useful suggestions and pointing me in the right directions in the world of CAT(0) cube complexes.

2 Background

Given a metric space X , we will always use $B_p(r)$ to denote the ball of radius r about the point $p \in X$.

Definition 2.1 (quasi-isometry). Let X and Y be metric spaces. A (k, c) -quasi-isometry is a not necessarily continuous map $f : X \rightarrow Y$, such that for all $a, b \in X$ we have:

$$\frac{1}{k}d_X(a, b) - c \leq d_Y(f(a), f(b)) \leq kd_X(a, b) + c$$

Quasi-isometries provide a natural notion of equivalence in a coarse geometric setting. For a detailed background on quasi-isometries and geometric group theory in general see [BH16].

2.1 Divergence

Let X be a metric space. Fix constants $0 < \delta \leq 1$, $\lambda \geq 0$ and consider the linear function $\rho(r) = \delta r - \lambda$. Let $a, b, c \in X$ and set $k = d(c, \{a, b\})$.

$$div_\gamma(a, b, c, \delta)$$

is the length of the shortest path in X from a to b which avoids the ball $B_c(\rho(k))$.

$$Div_\gamma^X(r, \delta)$$

is the supremum of $div_\gamma(a, b, c, \delta)$ over all a, b, c with $d(a, b) \leq r$. We will often suppress δ and γ from the notation and say the *divergence* of X is the function $Div(X) = Div_\gamma^X(r, \delta)$.

We set $f(r) \asymp g(r)$ if there exists a C such that:

$$\frac{1}{C}g\left(\frac{r}{C}\right) - Cr - C < f(r) < Cg(Cr) + Cr + C$$

Up to this equivalence relation on functions and under mild assumptions on the metric space, divergence is a quasi-isometry invariant. See Lemma 3.4 [DMS10] for the relevant hypotheses. For instance, the Cayley graph of a finitely generated group satisfies such hypotheses.

2.2 CAT(0) Cube Complexes

A *CAT(0) cube complex*, X , is a simply connected cell complex whose cells consist of Euclidean unit cubes, $[-\frac{1}{2}, \frac{1}{2}]^d$, of varying dimension d . Additionally, the link of each vertex is a flag complex (i.e., any set of vertices which are pairwise connected by an edge, span a simplex). X with the induced metric is a CAT(0) space (see [CS11] and [Wis11] for more detailed references). X is *finite-dimensional* if there is an upper bound on the dimension of cubes in X .

A *midcube* $Y \subset [-\frac{1}{2}, \frac{1}{2}]^d$ is the restriction of a coordinate to 0. A *hyperplane* $\mathcal{H} \subset X$ is the subspace with the property that for each cube C in X , $\mathcal{H} \cap C$ is a midcube or

$\mathcal{H} \cap C = \emptyset$. $X - \mathcal{H}$ consists of exactly two distinct components. A *half-space* is the closure of such a component. We denote the two half-spaces associated to \mathcal{H} by \mathcal{H}^+ and \mathcal{H}^- . The carrier of a hyperplane, $N(\mathcal{H})$, is the set of all cubes in X which have non-trivial intersection with \mathcal{H} .

In this paper, we will mostly work with the combinatorial metric on the 1-skeleton of X . A *combinatorial geodesic* is a geodesic in the 1-skeleton of X under this metric. The following core lemmas, whose proofs are found in [CS11], are used throughout this paper.

Lemma 2.2. *Suppose X is a finite-dimensional $CAT(0)$ cube complex. X is quasi-isometric to its 1-skeleton endowed with the combinatorial metric.*

Lemma 2.3. *Suppose X is a finite-dimensional $CAT(0)$ cube complex. For each $k > 0$, there exists a number $N(k)$ such that any combinatorial geodesics of length $N(k)$ in X must cross a set of pairwise non-intersecting hyperplanes $\{\mathcal{H}_1, \dots, \mathcal{H}_k\}$.*

2.3 Coxeter Groups

We only give a brief background on Coxeter groups. For an extensive background we refer the reader to [BB05] and [Dav08].

In this paper Γ will always denote a simplicial graph (usually a Coxeter graph). $V(\Gamma)$ and $E(\Gamma)$ are respectively the vertex set and edge set of Γ . A *clique* in Γ is a subgraph whose vertices are all pairwise adjacent. A k -clique is a clique with k vertices.

For $s \in \Gamma$, the *link of s* , $Link(s) \subset V(\Gamma)$, is the set of vertices in Γ connected to s by an edge. The *star of s* is the set $Star(s) = Link(s) \cup s$.

A Coxeter group is defined by the presentation:

$$W = \langle s_1, s_2, \dots, s_n \mid (s_i s_j)^{m(s_i, s_j)} = 1 \rangle$$

where $m(s_i, s_i) = 1$ and $m(s_i, s_j) = m(s_j, s_i) \in \{2, 3, \dots, \infty\}$ when $i \neq j$. If $m(s_i, s_j) = \infty$ then no relation of the form $(s_i s_j)^m$ is imposed.

Given a presentation for a Coxeter group, there is a corresponding labeled *Coxeter graph* Γ . The vertices of Γ are elements of S . There is an edge between s_i and s_j if and only if $m(s_i, s_j) \neq \infty$. This edge is labeled by $m(s_i, s_j)$ if $m(s_i, s_j) \geq 3$. If $m(s_i, s_j) = 2$ no label is placed on the corresponding edge. Conversely, given such an edge-labeled graph Γ , we have the Coxeter group W_Γ .

In the literature, there are many different conventions for associating a graph to a Coxeter group presentation. The given convention was chosen to make the theorems in this paper easier to state.

A *right-angled Coxeter group (RACG)* is a Coxeter group with generating set S where $m(s, t) \in \{\infty, 2\}$ for s, t distinct elements in S . An *even Coxeter group* is a Coxeter group given by a Coxeter graph where each edge either has an even label or no label.

We will often want to consider subgroups of a Coxeter group W_Γ corresponding to subgraphs of Γ . The full subgraph of $T \subset V(\Gamma)$ is the graph with vertex set T with a labeled edge (t_1, t_2) if and only if (t_1, t_2) is an edge of Γ with the same label.

Definition 2.4 (Induced Subgroup). Let (W, S) be a Coxeter group. For $T \subset S$, let W_T be the subgroup of W generated by the full subgraph of T . W_T is isomorphic to the Coxeter group corresponding to the subgraph of Γ induced by T (see [BB05] for instance).

The *Davis complex*, Σ_Γ , is a natural CAT(0) cell complex which the Coxeter group W_Γ acts geometrically. In this paper, we will only make use of the Davis complex for right-angled Coxeter groups.

Suppose W_Γ is a RACG. For every k -clique, $T \subset \Gamma$, the induced subgroup W_T is isomorphic to the direct product of k copies of \mathbb{Z}_2 . Hence, the Cayley graph of W_T is isomorphic to a k -cube. The Davis complex Σ_Γ has 1-skeleton the Cayley graph of W_Γ where edges are given unit length. Additionally, for each k -clique, $T \subset \Gamma$, and coset, gW_T , we glue a unit k -cube to $gW_T \subset \Sigma_\Gamma$. It is then evident that the Davis Complex for a RACG is naturally a CAT(0) cube complex.

3 Disk Diagrams in CAT(0) Cube Complexes

A *disk diagram* D is a contractible finite 2-dimensional cube complex with a fixed planar embedding $P : D \rightarrow \mathbb{R}^2$. The *area* of D is the number of 2-cells it contains. By compactifying \mathbb{R}^2 , $S^2 = \mathbb{R}^2 \cup \infty$, we can extend $P : D \rightarrow S^2$, giving a cellulation of S^2 . The *boundary path* of D , ∂D , is the attaching map of the cell in this cellulation containing ∞ . Note that this is not necessarily the topological boundary.

Let X be a CAT(0) cube complex. D is a *disk diagram in X* , if D is a disk diagram and there is a fixed continuous combinatorial map of cube complexes $F : D \rightarrow X$. By a lemma of Van Kampen, for every null-homotopic closed combinatorial path $p : S^1 \rightarrow X$, there exists a disk diagram D in X such that $\partial D = p$.

Suppose D is a disk diagram in a CAT(0) cube complex X and t is a 1-cell of D . A *dual curve H dual to t* , is a concatenation of midcubes in D which contains a midcube in D which intersects t . The image of H under the map $F : D \rightarrow X$ lies in some hyperplane $\mathcal{H} \subset X$. We also have that every edge in D is dual to exactly one maximal dual curve.

In our notation, we denote a dual curve by a capital letter and its corresponding hyperplane by the corresponding script letter.

An *end* of a dual curve H in D is a point of intersection of H with ∂D . Maximal dual curves either have no ends or two ends. The *carrier* $N(H)$ of H is the set of 2-cubes in D containing H .

Suppose we have oriented paths p_1, p_2, \dots, p_n in a disk diagram D and that $p_i \cap p_{i+1} \neq \emptyset$ for $1 \leq i < n$. We then define a new oriented path $p = p_1 * p_2 * \dots * p_n$ by beginning at the first point of p_1 , followed by p_1 until it's first intersection with p_2 ; followed by p_2 until its first intersection with p_3 . In this definition, we further assume the orientations are chosen such that this construction is possible.

Many of the ideas in this section originated in [Hag13a] and [Wis11]. For our purposes, we require some modified definitions and lemmas to those in the mentioned works. For completeness we include proofs of these modified claims, even though many of the arguments are very similar.

To every closed loop formed by a concatenation of combinatorial paths and hyperplanes, we wish to associate a disk diagram with boundary path this loop. This notion is formally defined below. If such a diagram is chosen appropriately, the dual curves associated to it behave nicely. We call such nicely behaved diagrams *combed* and define them later in this section.

Definition 3.1 (Hyperplane-Path Sequence). In the following definition, we work modulo $n + 1$ (i.e, $n + 1 = 0$). Let $\bar{A} = \{A_0, A_1, \dots, A_n\}$ be a sequence such that for

each i , A_i is either a hyperplane or an oriented combinatorial path in a CAT(0) cube complex X . Furthermore, for $0 \leq i \leq n$, if A_i and A_{i+1} are both hyperplanes then they intersect. If A_i and A_{i+1} are both combinatorial paths, then the endpoint of A_i is the initial point of A_{i+1} . If A_i is a hyperplane and A_{i+1} is a path, then the beginning point of A_{i+1} lies on $N(A_i)$. Similarly, if A_i is a path and A_{i+1} is a hyperplane, then the endpoint of A_i lies on $N(A_{i+1})$. We call \bar{A} a *hyperplane-path sequence*.

Given a hyperplane-path sequence $\bar{A} = \{A_0, \dots, A_n\}$, let $\bar{P} = \{P_0, \dots, P_n\}$ be a sequence of combinatorial paths where $P_i = A_i$ if A_i is a combinatorial path and P_i is a geodesic in $N(A_i)$ if A_i is a hyperplane. Furthermore, assume $P = P_0 * P_1 * \dots * P_n$ defines a loop. A disk diagram D is *supported by* \bar{A} if $\partial D = P$ for some choice of \bar{P} . Often this choice is given and we say D is *supported by* \bar{A} *with boundary path* \bar{P} . For an example of a disk diagram supported by a hyperplane path sequence see Figure 3.

Remark 3.1.1. Diagrams supported by a hyperplane-path sequence are a special case of diagrams with fixed carriers defined in [Hag13a]. The difference is that consecutive hyperplanes in a hyperplane-path sequence must intersect, where in [Hag13a] they either intersect or osculate. Most of this section can be modified to allow for osculating hyperplanes; however, there was no need for such sequences in this paper.

Definition 3.2 (Nongons, Bigons, Monogons and Oscugons). A *nongon* is a dual curve of length greater than one which begins and ends on the same dual 1-cell. A *bigon* is a pair of dual curves which intersect at their first and last containing squares. A *monogon* is a dual curve which intersects itself in its first and last square. An *oscugon* is a dual curve which starts at the dual 1-cell e_1 , ends at the dual 1-cell e_2 , such that $e_1 \neq e_2$, $e_1 \cap e_2 \neq \emptyset$ and e_1, e_2 are not contained in a common square. A disk diagram *without pathologies* is one that does not contain a nongon, bigon, monogon or oscugon.

The following is proven in [Wis11, Corollary 2.4]:

Lemma 3.3 ([Wis11]). *Suppose D is a disk diagram in a CAT(0) cube complex X . D does not contain monogons.*

We now wish to discuss the idea of *boundary combinatorics* in a disk diagram D . Suppose D and D' are two disk diagrams in a CAT(0) cube complex X . Let $p \subset \partial D$ and $p' \subset \partial D'$ be subcomplexes. p and p' are *equal boundary complexes* if the canonical maps $p \rightarrow X$ and $p' \rightarrow X$ are the same combinatorial maps.

Suppose $p \subset \partial D$ and $p' \subset \partial D'$ are equal boundary complexes, and let $i : p \rightarrow p'$ be the canonical isomorphism between them. p and p' have *equal boundary combinatorics* if given any pair of edges $e_1, e_2 \in p$ dual to a common dual curve in D , it follows $i(e_1)$ and $i(e_2)$ are dual to a common dual curve in D' .

In particular, D and D' have *equal boundary* if ∂D and $\partial D'$ are equal combinatorial complexes. Additionally, D and D' have *equal boundary combinatorics* if ∂D and $\partial D'$ have equal boundary combinatorics.

The following is also proven in [Wis11, Lemma 2.3]:

Lemma 3.4 ([Wis11]). *Let D be a disk diagram in a CAT(0) cube complex X . There exists a disk diagram D' in X with no pathologies which has equal boundary combinatorics to D .*

Definition 3.5 (Combed Diagram). Let D be a disk diagram supported by hyperplane-path sequence $\bar{A} = \{A_0, \dots, A_n\}$ with boundary path $\bar{P} = \{P_0, \dots, P_n\}$. We say D is *combed* if the following properties hold:

1. D has no pathologies.
2. If A_i is a hyperplane, no two dual curves dual to P_i intersect.
3. If A_i and A_{i+1} are both hyperplanes, no dual curve dual to P_i intersects P_{i+1} .

The arguments in the next two Lemmas are essentially the same as those in the proof of [Hag13a, Lemma 2.11]. However, for our purposes, we often require a statement regarding the boundary combinatorics of a given disk diagram. To be self-contained and to guarantee the arguments are valid in our context, we provide them here.

Lemma 3.6. *Suppose D is a disk diagram supported by hyperplane-path sequence $\bar{A} = \{A_0, \dots, A_n\}$ with boundary path $\bar{P} = \{P_0, \dots, P_n\}$, satisfying properties 1 and 2 of combed diagrams (Definition 3.5). There exists a combed diagram D' with boundary path $\bar{P}' = \{P'_0, \dots, P'_n\}$, where P'_i is a connected subsegment of P_i . Furthermore, \bar{P}' has equal boundary combinatorics as its image in \bar{P} .*

Proof. Suppose some dual curve C intersects both P_i and P_{i+1} . Let v be the vertex where P_i meets P_{i+1} . Let e_1 be the edge in P_i which intersects v and e_2 the edge in P_{i+1} which intersects v .

By property 2 of combed diagrams, every dual curve to P_i between v and C intersects P_{i+1} as well. It now follows that the dual curve to e_1 intersects e_2 . Since D contains no bigons, it follows $e_1 = e_2$. Hence, we obtain a new diagram from D by simply deleting the edge e_1 . By repeating this process we obtain the desired diagram D' . □

Lemma 3.7. *Let D be a disk diagram in a $CAT(0)$ cube complex supported by hyperplane-path sequence $\bar{A} = \{A_0, \dots, A_n\}$ with boundary path $\bar{P} = \{P_0, \dots, P_n\}$. Suppose for some i , $\mathcal{A} = A_i$ is a hyperplane. There exists a disk diagram D' also supported by $\bar{A} = \{A_0, \dots, A_n\}$ with boundary path $\bar{P}' = \{P_0, \dots, P_{i-1}, P'_i, P_{i+1}, \dots, P_n\}$ such that $\partial D - P_i$ has the same boundary combinatorics as $\partial D' - P'_i$. Additionally, no two dual curves dual to P'_i in D' intersect.*

Proof. By Lemma 3.4 we may assume D has no pathologies. Set $P = P_i$ and suppose two dual curves to P , C_1 and C_2 , intersect. If $d_P(N(C_1), N(C_2)) = d > 0$, it follows there is another dual curve to P , C_3 , between C_1 and C_2 in P . C_3 must then intersect either C_1 or C_2 . Without loss of generality, assume C_3 intersects C_1 and note that $d_P(N(C_1), N(C_3)) < d$. Hence, proceeding this way, and under the assumption of no bigons, we may deduce there is a square S in D' with two of its edges lying on P .

Label the edges of S as $\{e_1, e_2, e_3, e_4\}$ and assume e_3 and e_4 are adjacent and lie in P . Let \mathcal{H}_1 and \mathcal{H}_2 respectively be hyperplanes which pass through e_1 and e_2 .

If $\mathcal{A} = \mathcal{H}_1$, then it follows that $e_1, e_2, e_3 \in N(\mathcal{A})$. Hence, $S \in N(\mathcal{A})$. Alternatively, suppose \mathcal{H}_1 , \mathcal{H}_2 , and \mathcal{A} are distinct. Since the link of vertices in X are flag complexes, it follows S is in a 3-cube which is contained in $N(\mathcal{A})$. In either case, we have that $S \in N(\mathcal{A})$.

Let P' be the path in D which is the same as P with e_3, e_4 replaced with e_1, e_2 . $|P'| = |P|$, so P' is geodesic. Furthermore, $P' \in N(\mathcal{A})$. Let $D' \subset D$ be the disk diagram obtained as a subdiagram of D by replacing P with P' . D' is a diagram of smaller area than D . Also note that the boundary combinatorics of $\partial D - P$ are not affected. Therefore, by iterating this process, we eventually have a diagram with the desired properties. □

The following Lemma guarantees the existence of a combed diagram.

Lemma 3.8. *Let D be a disk diagram supported by hyperplane-path sequence \bar{A} . There exists a combed disk diagram D' also supported by \bar{A} .*

Proof. The Lemma follows by applying Lemma 3.4 to D to get a disk diagram with no pathologies. We then repeatedly apply Lemma 3.7 to the resulting diagram to obtain a diagram satisfying property 1 and 2 of combed diagrams. Finally, we apply Lemma 3.6 to obtain a combed diagram. \square

Given a maximal dual curve C in a disk diagram D , we want to construct a new combed diagram with the hyperplane C in its support. Furthermore, we want the appropriate boundary combinatorics of D preserved in this new diagram. The following technical Lemma guarantees the existence of such a diagram.

Lemma 3.9. *Let D be a combed disk diagram in a $CAT(0)$ cube complex X supported by hyperplane-path sequence $\bar{A} = \{A_0, \dots, A_n\}$ and with boundary path $\bar{P} = \{P_0, \dots, P_n\}$.*

Suppose C is a maximal dual curve from P_i to P_j with $i < j$ and let $\mathcal{C} \subset X$ be the hyperplane associated to C . Let P be a combinatorial path in $N(\mathcal{C})$ from P_i to P_j . Let P'_i be the subsegment of P_i between P_{i-1} and P , and let P'_j be the subsegment of P_j between P and P_{j+1} . Let A'_i and A'_j be the corresponding supports of P'_i and P'_j respectively ($A'_i = A_i$ if A_i is a hyperplane and $A_i = P'_i$ otherwise).

There exists a combed disk diagram D' supported by

$$\bar{A}' = \{A_0, \dots, A_{i-1}, A'_i, \mathcal{C}, A'_j, A_{j+1}, \dots, A_n\}$$

with boundary path

$$\bar{P}' = \{P_0, \dots, P_{i-1}, P'_i, P', P'_j, P_{j+1}, \dots, P_n\}$$

such that $\partial D' - P'$ has the same boundary combinatorics as the corresponding subset of ∂D .

Proof. Let P' be a geodesic in \mathcal{C} connecting the endpoints of P . Let D_1 be a combed diagram with boundary path $\{P', P\}$. Let D_2 be the subdiagram of D with boundary path $\{P_0, \dots, P'_i, P, P'_j, \dots, P_n\}$. Note that D_2 is still combed. We may form a new disk diagram D_3 by gluing D_1 to D_2 along P . D_3 is supported by $\{A_0, \dots, A'_i, \mathcal{C}, A'_j, \dots, A_n\}$ with boundary path $\{P_0, \dots, P'_i, P', P'_j, \dots, P_n\}$.

By Lemma 3.4, we may assume D_3 has no pathologies and $\partial D_3 - P'$ has the same combinatorics as the corresponding subset of ∂D .

All that is left to prove is that properties 2 and 3 of combed diagrams (definition 3.5) hold. Property 3 clearly still holds for dual curves which do not intersect P' . Assume A_i is a hyperplane. Since D is combed, no dual curve to P'_i in D_2 intersects P . Since the combinatorics of D_2 are preserved in D_3 , this is still the case in D_3 . So, in particular, no dual curve to P'_i intersects P' in D_3 . Therefore, property 3 holds in D_3 .

Assume property 2 is false in D_3 . We then have that two dual curves C_1, C_2 , dual to the same boundary path in D_3 with hyperplane support, intersect. By the preservation of boundary combinatorics and the fact that D_3 is combed, it follows each of these curves must have an endpoint on P' . We can then modify D_3 using Lemma 3.7 to obtain a new diagram D' satisfying the claim. \square

4 Hyperplane Separation Properties

For the remainder of the paper, X will denote a CAT(0) cube complex. Furthermore, \mathcal{Y} and \mathcal{Z} will always denote a pair of non-intersecting unbounded hyperplanes in X . If X is one-ended and essential then every hyperplane is unbounded.

We will discuss different definitions for separation properties of a given pair of non-intersecting hyperplanes. In the next section we will explore the relationship between these separation properties and the divergence of X .

Definition 4.1 and Lemma 4.2 given below are from [BC12].

Definition 4.1 ([BC12]). \mathcal{Y} and \mathcal{Z} are *strongly separated* if no hyperplane intersects them both.

A *minimal geodesic* g between hyperplanes \mathcal{Y} and \mathcal{Z} is a geodesic with endpoints on \mathcal{Y} and \mathcal{Z} such that $|g|$ is minimal over all such geodesics.

Lemma 4.2 ([BC12]). *There exists only one minimal geodesic connecting a pair of strongly separated hyperplanes.*

The following definition gives a slight generalization of strongly separated hyperplanes.

Definition 4.3. \mathcal{Y} and \mathcal{Z} are k -separated if at most k hyperplanes intersect both \mathcal{Y} and \mathcal{Z} .

In particular, a pair of strongly separated hyperplanes are 0-separated. The following Lemma describes how the set of geodesics and hyperplanes intersecting k -separated hyperplanes behave nicely. It should be compared to Lemma 4.2.

Lemma 4.4. *Suppose \mathcal{Y} and \mathcal{Z} are k -separated, then $d(\mathcal{H}_1 \cap \mathcal{Y}, \mathcal{H}_2 \cap \mathcal{Y}) \leq k$ for every pair of hyperplanes $\mathcal{H}_1, \mathcal{H}_2$ which intersect both \mathcal{Y} and \mathcal{Z} . Furthermore, every minimal geodesic g connecting \mathcal{Y} to \mathcal{Z} either lies in the carrier of a hyperplane which intersects both \mathcal{Y} and \mathcal{Z} , or alternatively, g is unique and \mathcal{Y} and \mathcal{Z} are strongly separated.*

Proof. Suppose hyperplanes \mathcal{H}_1 and \mathcal{H}_2 each intersect both \mathcal{Y} and \mathcal{Z} . For a contradiction, suppose that $d(\mathcal{H}_1 \cap \mathcal{Y}, \mathcal{H}_2 \cap \mathcal{Y}) > k$. Let D be a combed disk diagram supported by $\{\mathcal{Y}, \mathcal{H}_1, \mathcal{Z}, \mathcal{H}_2\}$. Every dual curve to \mathcal{Y} must intersect \mathcal{Z} . However, there are more than k such dual curves and hence more than k hyperplanes intersecting both \mathcal{Y} and \mathcal{Z} . This contradicts \mathcal{Y} and \mathcal{Z} being k -separated.

To prove the Lemma's second claim, suppose g is a geodesic from \mathcal{Y} to \mathcal{Z} and that g does not lie in a hyperplane. If \mathcal{Y} and \mathcal{Z} are strongly separated, then by Lemma 4.2, g is unique. Otherwise, suppose \mathcal{H} is a hyperplane intersecting both \mathcal{Y} and \mathcal{Z} and let D be a combed diagram supported by the hyperplane-path sequence $\{\mathcal{Y}, g, \mathcal{Z}, \mathcal{H}\}$ with boundary path $\{Y, g, Z, H\}$. Every dual curve to H must intersect g . Since g is geodesic, $|g| = |H|$ and it follows no dual curve to \mathcal{Y} can intersect g . So every dual curve to Y intersects Z . It follows D is an Euclidean rectangle. Hence, there is a dual curve C from Y to Z which has g as part of its boundary path. So, g is contained in the carrier of the hyperplane \mathcal{C} which intersects both \mathcal{Y} and \mathcal{Z} . \square

Definition 4.5. If infinitely many hyperplanes intersect both \mathcal{Y} and \mathcal{Z} then we say \mathcal{Y} and \mathcal{Z} are ∞ -connected.

A pair of non-intersecting hyperplanes are k -chain connected (formally defined below) if there is an appropriate sequence of sets of hyperplanes connecting them. k -chain separated hyperplanes (hyperplanes which are not k -chain connected) provide a generalization of the notion of k -separated hyperplanes.




Figure 1: The hyperplanes \mathcal{Y} and \mathcal{Z} above are 3-chain connected.

Definition 4.6 (k -chain connected). \mathcal{Y} and \mathcal{Z} are k -chain connected if there exists a sequence of length k sequences of hyperplanes:

$$S_1 = \{\mathcal{H}_1^1, \mathcal{H}_2^1, \dots, \mathcal{H}_k^1\}$$

$$S_2 = \{\mathcal{H}_1^2, \mathcal{H}_2^2, \dots, \mathcal{H}_k^2\}$$

...

$$S_m = \{\mathcal{H}_1^m, \mathcal{H}_2^m, \dots, \mathcal{H}_k^m\}$$

satisfying the following properties:

- I. For each i , hyperplanes in S_i pairwise do not intersect.
- II. For each $i < m$, each hyperplane in S_i intersects each hyperplane in S_{i+1} .
- III. Every hyperplane in S_1 intersects \mathcal{Y} and every hyperplane in S_m intersects \mathcal{Z} .

Definition 4.7. \mathcal{Y} and \mathcal{Z} are k -chain separated if they are not k -chain connected.

Definition 4.8. Hyperplanes \mathcal{H} and \mathcal{H}' are of the same type, if they are in the same orbit of $\text{Aut}(X)$. \mathcal{H} and \mathcal{H}' are of non-intersecting type if $g\mathcal{H} \cap \mathcal{H}' = \emptyset$ for all $g \in \text{Aut}(X)$.

Definition 4.9. Let $Q = \{\mathcal{H}_1, \dots, \mathcal{H}_m\}$ be a sequence of hyperplanes. Define

$$\text{Type}(Q) = (T_1, \dots, T_m)$$

where T_i is the hyperplane type (orbit class) of \mathcal{H}_i . Note that the tuple $\text{Type}(Q)$ is ordered.

The next set of definitions provide a further strengthening of the notion of k -chain separated hyperplanes which allows us to prove stronger divergence bounds in the next section. The following definitions were created with the key example of right-angled Coxeter groups in mind.




Figure 2: Hyperplanes \mathcal{Y} and \mathcal{Z} are symbolically 2-chain connected. The hyperplane colors signify their type.

Definition 4.10 (Symbolically k -chain Connected). \mathcal{Y} and \mathcal{Z} are *symbolically k -chain connected* if there exists a sequence of length k sequences of hyperplanes:

$$S_1 = \{\mathcal{H}_1^1, \mathcal{H}_2^1, \dots, \mathcal{H}_k^1\}$$

$$S_2 = \{\mathcal{H}_1^2, \mathcal{H}_2^2, \dots, \mathcal{H}_k^2\}$$

...

$$S_m = \{\mathcal{H}_1^m, \mathcal{H}_2^m, \dots, \mathcal{H}_k^m\}$$

satisfying the following five properties:

- I. For each $i \leq m$ and $j < k$, \mathcal{H}_j^i and \mathcal{H}_{j+1}^i are of non-intersecting type.
 - II. For each $1 < i \leq m$, every hyperplane in S_i intersects \mathcal{H}_1^{i-1} .
 - III. For each $i \leq m$ and $j \leq k$, \mathcal{H}_j^i intersects every hyperplane in $S_{i'}$ for some $i' > i$.
- Define the function $F(\mathcal{H}_j^i) = \text{Type}(S_{i'})$. Additionally, for all $j, j' \leq k$ and for all $i < m$, $F(\mathcal{H}_j^i) = F(\mathcal{H}_{j'}^i)$.

- IV. Every hyperplane in S_1 intersects \mathcal{Y} and every hyperplane in S_m intersects \mathcal{Z} .
- V. Let g be a minimal geodesic from \mathcal{Y} to \mathcal{Z} , for all $i \leq m$ and $j \leq k$, g and \mathcal{H}_{j-1}^i lie in different half-spaces of \mathcal{H}_j^i .

Remark 4.10.1. By Lemma 4.4, property V in the definition above necessarily implies that $m > 1$. Furthermore, it follows that if property V is true for a geodesic from \mathcal{Y} to \mathcal{Z} , then it is true for all geodesics from \mathcal{Y} to \mathcal{Z} .

Definition 4.11 (symbolically k -chain separated). Two non-intersecting hyperplanes are *symbolically k -chain separated* if they are not symbolically k -chain connected and are not k -connected.

5 Divergence in CAT(0) Cube Complexes

We now define the *hyperplane divergence function*, $HDiv$, which measures the length of a shortest path between two hyperplanes which avoids a ball centered on one of the hyperplanes. Later in this section, we obtain lower bounds on the Div function from lower bounds on the $HDiv$ function.

Definition 5.1. Let g be a minimal geodesic from \mathcal{Y} to \mathcal{Z} and set $p = \mathcal{Y} \cap g$. $HDiv_g(\mathcal{Y}, \mathcal{Z})(r)$ is the length of a shortest path from \mathcal{Y} to \mathcal{Z} which avoids the ball $B_p(r)$.

Remark 5.1.1. If X is one-ended then $HDiv_g(\mathcal{Y}, \mathcal{Z})(r)$ always takes finite values. Additionally, if \mathcal{Y} and \mathcal{Z} are k -separated and g_1, g_2 are different minimal geodesics from \mathcal{Y} to \mathcal{Z} , then by Lemma 4.4 we have that $HDiv_{g_1}(\mathcal{Y}, \mathcal{Z})(r - k) \leq HDiv_{g_2}(\mathcal{Y}, \mathcal{Z})(r) \leq HDiv_{g_1}(\mathcal{Y}, \mathcal{Z})(r + k)$. Hence, up to the usual equivalence on divergence functions, for k -separated hyperplanes it is often not relevant which minimal geodesic is used.

This section is devoted to proving the following two theorems which provide a connection between the hyperplane separation properties defined in the previous section and divergence in X . Theorem 5.2 gives bounds on $HDiv(\mathcal{Y}, \mathcal{Z})$, and Theorem 5.3 gives bounds on $Div(X)$.

Theorem 5.2. *The following are true:*

1. *Suppose X is finite-dimensional and locally compact. \mathcal{Y} and \mathcal{Z} are ∞ -connected if and only if $HDiv(\mathcal{Y}, \mathcal{Z})$ is constant.*
2. *If \mathcal{Y} and \mathcal{Z} are k -separated, then $HDiv(\mathcal{Y}, \mathcal{Z})$ is at least linear.*
3. *If \mathcal{Y} and \mathcal{Z} are k -chain separated and X is finite-dimensional, then $HDiv(\mathcal{Y}, \mathcal{Z})(R) \succeq \frac{1}{2} \log_2(\log_2 R)R$.*

Theorem 5.3. *Suppose X is essential, locally compact and with cocompact automorphism group.*

1. *If \mathcal{Y} and \mathcal{Z} are k -separated, then $Div(X)$ is bounded below by a quadratic function.*
2. *If \mathcal{Y} and \mathcal{Z} are k -chain separated, then $Div(X) \succeq \frac{1}{2} \log_2(\log_2(R))R^2$.*
3. *Suppose X has k -alternating geodesics (Definition 5.4). If \mathcal{Y} and \mathcal{Z} are symbolically k -chain separated then $HDiv(\mathcal{Y}, \mathcal{Z})$ is bounded below by a quadratic function and $Div(X)$ is bounded below by a cubic function.*

Definition 5.4. X has k -alternating geodesics, if there exists a constant M so that every geodesic of length M in X crosses a set of hyperplanes $\{\mathcal{H}_1, \dots, \mathcal{H}_k\}$ such that \mathcal{H}_i and \mathcal{H}_{i+1} are of non-intersecting type for all $i < k$.

Lemma 5.5. Suppose X is finite-dimensional, locally compact and that \mathcal{Y} and \mathcal{Z} are ∞ -connected. There exists a constant c such that for all r and choice of geodesic g , $HDiv_g(\mathcal{Y}, \mathcal{Z})(r) = c$.

Proof. Fix a hyperplane H_1 intersecting both \mathcal{Y} to \mathcal{Z} and let \mathcal{H}_2 be another hyperplane intersecting both \mathcal{Y} to \mathcal{Z} a distance at least r from \mathcal{H}_1 . Let D be a combed disk diagram supported by $\{\mathcal{Y}, \mathcal{H}_1, \mathcal{Z}, \mathcal{H}_2\}$. Every dual curve to \mathcal{H}_1 must intersect \mathcal{H}_2 and every dual curve to \mathcal{Y} must intersect \mathcal{Z} . Hence, D is an Euclidean strip of dimension $r \times d(\mathcal{Y}, \mathcal{Z})$ and the lemma follows. \square

Lemma 5.6. Suppose \mathcal{Y} and \mathcal{Z} are k -separated. There exists a constant c such that $HDiv(\mathcal{Y}, \mathcal{Z})(r) \geq r + c$.

Proof. Let g be any minimal geodesic from \mathcal{Y} to \mathcal{Z} and $p = g \cap \mathcal{Y}$. Let α be a path from \mathcal{Y} to \mathcal{Z} which avoids the ball $B_p(r)$. Let D be a combed disk diagram supported by $\{\mathcal{Y}, \alpha, \mathcal{Z}, g^{-1}\}$. At most $d = |g|$ dual curves to \mathcal{Y} can intersect g and at most k dual curves to \mathcal{Y} can intersect \mathcal{Z} . Hence, at least $r - d - k$ dual curves to \mathcal{Y} intersect α . Hence, $|\alpha| \geq r - d - k$. \square

Lemma 5.7. Suppose X is finite-dimensional and \mathcal{Y}, \mathcal{Z} are k -chain separated. Set $d = d_X(\mathcal{Y}, \mathcal{Z})$. There exists a constant $R_0(d, k)$ such that for $R > R_0$, $HDiv(\mathcal{Y}, \mathcal{Z}) \geq \frac{1}{2}R \log_2(\log_2 R)$.




Figure 3: Graphic for the proof Lemma 5.7. D_3 is a diagram supported by the hyperplane-path sequence $\{\mathcal{H}_0, \mathcal{H}_1, \mathcal{H}_2, \mathcal{H}_3, \alpha_3, \mathcal{Z}, g^{-1}\}$.

Proof. By Lemma 2.3, let $K > 0$ be the constant, only depending on k and X , such that a geodesic of length K in X must intersect at least k pairwise non-intersecting hyperplanes. Let g be a minimal geodesic from \mathcal{Y} to \mathcal{Z} and set $p = g \cap \mathcal{Y}$. Fix $R > 0$

and let $r = \log_2(\log_2 R)$. Let α be a combinatorial path from \mathcal{Y} to \mathcal{Z} which avoids the ball $B_p(R)$.

Set $\mathcal{H}_0 = \mathcal{Y}$. Let D_0 be a combed disk diagram with boundary supported by the hyperplane-path sequence $\bar{A}_0 = \{\mathcal{H}_0, \alpha, \mathcal{Z}, g^{-1}\}$. Orient H_0 from g to α . For $i \in \mathbb{Z}_{\geq 0}$ set $c_i = 2^{i+1}r(K+d)$ where $d = |g|$.

For $n \leq r$, define inductively H_n as the c_{n-1} 'th dual curve to H_{n-1} in D_{n-1} and assume H_n intersects α . Orient H_n from H_{n-1} to α . Define α_n as the subpath of α from H_n to Z and β_n as the subpath of α from \mathcal{H}_{n-1} to \mathcal{H}_n . Define D_n as the combed diagram with boundary supported by

$$\bar{A}_n = \{\mathcal{H}_0, \mathcal{H}_1, \dots, \mathcal{H}_n, \alpha_n, \mathcal{Z}, g^{-1}\}$$

obtained from D_{n-1} by Lemma 3.9. For $j \leq n$, define the paths in D_n :

$$T_j = H_{j+1} * H_{j+2} * \dots * H_n * \alpha_n \text{ (set } T_n = \alpha_n)$$

$$B_j = Z^{-1} * g^{-1} * H_0 * H_1 * \dots * H_{j-1}$$

$$B'_j = g^{-1} * H_0 * H_1 * \dots * H_{j-1}$$

We will show through induction the following are true for all $n \leq r$:

- A. D_n is well defined.
- B. For $j < n$, rK dual curves to H_j in D_n , intersect T_j .
- C. For $n > 0$, $|\beta_n| \geq R - c_n$

Given the diagram D_{n-1} , in order to define D_n we must first show H_{n-1} has at least c_{n-1} dual curves emanating from it in D_{n-1} . Since D_n only needs to be defined for $n \leq r$, we can do this by showing $|H_r| > 0$ in D_r . Because α avoids the R -ball about p , we have:

$$\begin{aligned} |H_r| &\geq R - \sum_{j=0}^{r-1} c_j \\ &= R - \sum_{j=0}^{r-1} 2^{j+1}r(K+d) = R - 2r(K+d) \sum_{j=0}^{r-1} 2^j \end{aligned}$$

Using the formula for a geometric series:

$$\begin{aligned} |H_r| &\geq R - 2r(K+d)(2^r - 1) \\ &= R - \left(2(K+d) \log_2(\log_2 R)\right) \left(2 \log_2 R - 1\right) \end{aligned}$$

There then exists a constant $m_1(k, d)$, such that for $R > m_1$, we have that $|H_r| \geq 0$. Thus when $R > m_1$, $H_n \geq c_n$ for $n < r$.

We now turn to the base case, $n = 0$. Hypothesis A. follows immediately. Let Q denote the set of dual curves to H_0 . At most d of these can intersect g . Additionally, we cannot have K dual curves in Q intersect Z . For if they did, this would imply \mathcal{Y} and \mathcal{Z} are k -chain connected, a contradiction. We then have that $R - K - d$ dual curves to H_0 intersect $\alpha = T_0 \subset D_0$.

By the following inequalities:

$$R > c_0 = 2r(K + d) > rK + K + d$$

hypothesis B. is shown true, settling the base case.

For the general case, assume $n + 1 \leq r$ and that hypotheses A-C are satisfied for any $n' < n + 1$. Let $Q = \{Q_1, Q_2, \dots, Q_m\}$ be the set of dual curves in D_n emanating from H_n ordered by the orientation on H_n . It follows at most $C = \sum_{j=0}^{n-2} c_j + d$ paths in Q can intersect B'_n (the sum does not go to $n - 1$ since the diagram is combed). Using the formula for a geometric series we have:

$$C = c_{n-1} - 2r(K + d) + d \leq c_{n-1}$$

Additionally, we cannot have K curves in Q intersect Z . For then, there is a subset of k of these dual curves, $S_1 = \{H_1^1, \dots, H_k^1\} \subset Q$, corresponding to pairwise non-intersecting hyperplanes, which intersect Z . By induction hypothesis B, k dual curves to H_{n-1} , $S_2 = \{H_1^2, H_2^2, \dots, H_k^2\}$, corresponding to pairwise non-intersecting hyperplanes, intersect every curve in S_1 . Now, $H_k^1 * H_k^2$ is a path from H_{n-1} to Z . By the induction hypothesis B and the pigeonhole principle, k dual curves emanating from H_{n-2} , $S_3 = \{H_1^3, \dots, H_k^3\}$, intersect either H_k^1 or H_k^2 . Hence, every curve in S_3 intersects every curve in S_1 or S_2 . Proceeding this way we can show \mathcal{Y} is k -chain connected to Z , a contradiction.

It follows for $j \geq c_{n-1} + K + d \geq C + K + d$, Q_j must intersect α . In particular, $H_{n+1} = Q_{c_n}$ must intersect α . Hence, using Lemma 3.9 we can define D_{n+1} , proving hypothesis A.

Note that for j such that, $c_{n-1} + K + d \leq j \leq c_n$, Q_j must intersect α . A direct calculation gives that there are at least rK of such curves. Because of this, and the boundary combinatorics preservation property of Lemma 3.9, hypothesis B is satisfied in D_{n+1} .

We are left to prove hypothesis C. Note that for $j > c_n$, Q_j intersects β_{n+1} in D_n . By using the fact that α avoids $B_p(R)$ we have:

$$\begin{aligned} |\beta_{n+1}| &\geq R - \sum_{j=0}^n c_j = R - 2r(K + d) \sum_{j=0}^n 2^j \\ &= R - 2r(K + d)(2^{n+1} - 1) = R - c_{n+1} + 2r(K + d) \\ &> R - c_{n+1} \end{aligned}$$

This proves induction hypothesis C and completes the induction.

We have thus divided α into a set of disjoint subpaths $\{\beta_i\}$ for each of which we have a length lower bound. This allows us to compute a lower bound for the length of α :

$$\begin{aligned} |\alpha| &\geq \sum_{i=1}^r \beta_i = \sum_{i=1}^r (R - c_i) \\ &= rR - \sum_{i=0}^{r-1} c_{i+1} \end{aligned}$$

$$\begin{aligned}
&= rR - 4r(K + d) \sum_{i=0}^{r-1} 2^i \\
&= rR - 4r(K + d)(2^r - 1) \\
&= \log_2(\log_2(R))R - 4(K + d)\log_2(\log_2 R) \left(\log_2 R - 1 \right)
\end{aligned}$$

There then exists a constant $m_2(k, d)$ such that for $R > m_2$

$$|\alpha| \geq \frac{\log_2(\log_2(R))R}{2}$$

□

We are now in a position to prove Theorem 5.2.

Proof of Theorem 5.2 . The theorem follows from the above three Lemmas.

□

The proof of the following Lemma is similar to that of Lemma 5.7. However, to get the quadratic bound on the $HDiv$ function the proof requires a different counting technique.

Lemma 5.8. *Suppose X is essential, locally compact, has k -alternating geodesics and that $Aut(X)$ acts cocompactly. Let \mathcal{Y} and \mathcal{Z} be symbolically k -chain separated. Set $d = d_X(\mathcal{Y}, \mathcal{Z})$. There exists a constant $R_0(d, k)$ such that for $R > R_0$, $HDiv(\mathcal{Y}, \mathcal{Z})$ is bounded below by a quadratic function.*

Proof. Fix $R > 0$. Let g be a minimal geodesic from \mathcal{Y} to \mathcal{Z} , $p = g \cap \mathcal{Y}$, and α a $B_p(R)$ avoidant combinatorial path from \mathcal{Y} to \mathcal{Z} . Let M be the k -alternating constant from Definition 5.4. Set $c_1 = M + d$. Let c_2 be the number of different hyperplane types in X (this is finite since $Aut(X)$ acts cocompactly). Set $c = c_1(c_2)^k + 2d$. Set $r = \frac{R}{6c}$ and set $\mathcal{H}_0 = \mathcal{Y}$. Let D_0 be a combed diagram with boundary supported by $\bar{A}_0 = \{\mathcal{Y}, \alpha, \mathcal{Z}, w_0^{-1}\}$ and orient H_0 from g to α .

Assume we have a combed disk diagram D_n supported by

$$\bar{A}_n = \{\mathcal{H}_0, \mathcal{H}_1, \dots, \mathcal{H}_n, \alpha_n, \mathcal{Z}, g^{-1}\}$$

where α_n is a subpath of α from \mathcal{H}_n to \mathcal{Z} . Assume H_n is oriented from H_{n-1} to α_n . Define β_n as the subpath of α from \mathcal{H}_{n-1} to \mathcal{H}_n .

Let $Q = \{Q_1, Q_2, \dots, Q_l\}$ be the set of dual curves to H_n labeled sequentially by the orientation on H_n . Define τ_n to be the largest integer such that Q_{τ_n} does not intersect α . Set $m_n = \tau_n + c$, and let $H_{n+1} = Q_{m_n}$. For $j \leq n$, define the paths in D_n :

$$T_j = H_{j+1} * H_{j+2} * \dots * H_n * \alpha_n \text{ (set } T_n = \alpha_n \text{)}$$

$$B_j = Z^{-1} * g^{-1} * H_0 * H_1 * \dots * H_{j-1}$$

$$B'_j = g^{-1} * H_0 * H_1 * \dots * H_{j-1} \subset B_j$$

We assume by induction the following are true for $n < r$:

A. D_n is defined.

- B.** In D_n , for $j < n$, exactly c dual curves to H_j intersect T_j and exactly τ_j dual curves to H_j intersect B_j .
- C.** For $n > 0$, $|\beta_n| \geq |H_{n-1}| - \tau_{n-1} - c$

For the case when $n = 0$, **A** follows immediately and **B**, **C** are trivial. Note that at most d curves can intersect g and M curves can intersect Z (since \mathcal{Y} and \mathcal{Z} are symbolically k -chain separated). Hence, $\tau_0 \leq d + M$, and Q_{m_0} does intersect α . Hence, by using Lemma 3.9, we can define D_1 . It is also clear that **B** holds in D_1 . Furthermore, in D_0 , it is clear that for $j > m_0$, Q_j intersects B_1 . Hence, **C** is true as well. This shows the base cases $n = 0$ and $n = 1$ are true.

We now turn to the general case. Suppose the lemma is true for all integers n' such that $n' \leq n < r$. Consider the diagram D_n , and let $Q = \{Q_1, \dots, Q_l\}$ be the set of dual curves to H_n labeled sequentially by the orientation on H_n .

Sub-Claim 5.9. *We will first show that we cannot have c_1 curves in Q intersect Z .*

Proof. We say two hyperplanes are almost symbolically k -chain connected, if they satisfy every condition of definition 4.10 except maybe condition V. Suppose for a contradiction the claim is not true. It follows that \mathcal{H}_n is almost symbolically k -chain connected to \mathcal{Z} . Assume for some $i \leq n$, \mathcal{H}_i and \mathcal{Z} are almost symbolically k -chain connected by sequences:

$$\begin{aligned} S_1 &= \{\mathcal{P}_1^1, \mathcal{P}_2^1, \dots, \mathcal{P}_k^1\} \\ S_2 &= \{\mathcal{P}_1^2, \mathcal{P}_2^2, \dots, \mathcal{P}_k^2\} \\ &\dots \\ S_m &= \{\mathcal{P}_1^m, \mathcal{P}_2^m, \dots, \mathcal{P}_k^m\} \end{aligned}$$

Additionally, we want this structure to be seen in the disk diagram D_n . So assume for every \mathcal{P}_i^j , there is a corresponding dual curve P_i^j in D_n . Also assume every dual curve corresponding to a hyperplane in S_1 intersects H_i , every dual curve corresponding to a hyperplane in S_m intersects Z , and $P_1^1 * P_1^2 * \dots * P_1^m$ is well defined as a path in D_n from H_i to Z .

By the induction hypothesis, there are c dual curves to H_{i-1} which intersect T_j . Let p be one such curve. Since D_n is combed, p cannot intersect T_{j+1} . Hence, p must intersect $P_1^1 * P_1^2 * \dots * P_1^m$, and, consequently p must intersect P_1^j for some j . However, since the curves in S_j are pairwise disjoint, p must intersect every curve in S_j .

There are only c_2^k different possibilities for the tuple $Type(S_j)$. Hence, by the pigeonhole principle, there must be k dual curves to H_{n-1} , $S = \{P_1, \dots, P_k\}$, such that the corresponding hyperplanes to the sequence:

$$\begin{aligned} S &= \{P_1, P_2, \dots, P_k\} \\ S_j &= \{P_1^1, P_2^1, \dots, P_k^1\} \\ S_{j+1} &= \{P_1^2, P_2^2, \dots, P_k^2\} \\ &\dots \\ S_m &= \{P_1^m, P_2^m, \dots, P_k^m\} \end{aligned}$$

almost symbolically k -chain connect \mathcal{H}_{n-1} to \mathcal{Z} .

Proceeding in this manner, this would imply \mathcal{Y} is symbolically k -chain connected to \mathcal{Z} , a contradiction. This finishes the proof of the subclaim. \square

We next want to show that m_n is well defined. By induction hypothesis **B** and the subclaim, at most $C = (n-2)c + d + c$ curves in Q can intersect B_n . Note that $\frac{R}{6} = rc \geq nc + c \geq C + c$. Hence, for m_n to be well defined, it is enough to know that $|H_n| \geq \frac{R}{6}$.

$\sum_{j=0}^{n-1} \tau_j$ is a count of how many dual curves have both endpoints on B_n . At most $(n-1)c$ such curves have endpoints on H_i and H_j for some $i < j < n$. At most d such curves have endpoints on g . Furthermore, by the subclaim at most $(n-1)c_1$ such curves have endpoints on H_i and Z for some $i < n$. Hence,

$$\sum_{j=0}^{n-1} \tau_j \leq (n-1)c + d + (n-1)c_1 \leq 2nc$$

Since α does not intersect $B_p(R)$, we have that:

$$\begin{aligned} |H_n| &\geq R - \sum_{i=0}^{n-1} |H_i| \\ &= R - \sum_{i=0}^{n-1} (\tau_i + c) = R - nc - \sum_{i=0}^{n-1} \tau_i \\ &\geq R - 3nc \\ &\geq R - 3rc = \frac{R}{2} \end{aligned}$$

Thus, m_n is well defined.

We are left to prove induction hypothesis **C**. For $j > m_n$, Q_j intersects β_{n+1} . Thus,

$$|\beta_n| \geq |H_{n-1}| - \tau_{n-1} - c$$

This finishes the induction.

We have broken α into a union of r subpaths $\{\beta_j\}$ for which we have a lower bound for. We can then calculate a lower bound for α :

$$\begin{aligned} |\alpha| &\geq \sum_{i=0}^{r-1} |\beta_{n+1}| \geq \sum_{i=0}^{r-1} |H_n| - \tau_n - c \\ &\geq \sum_{i=0}^{r-1} \left(\frac{R}{2} - \tau_i - c \right) \\ &\geq \frac{rR}{2} - \sum_{i=0}^{r-1} (\tau_i) - rc \\ &\geq \frac{rR}{2} - 3rc = \frac{R^2}{12c} - \frac{R}{2} \end{aligned}$$

\square

The following theorem allows us to deduce stronger lower bounds for $Div(X)$ through the existence of just two hyperplanes with strong enough separation properties. The proof of the theorem involves constructing an infinite sequence of nested hyperplanes. This is done primarily through the machinery developed in [CS11].

Theorem 5.10. *Let X be essential, locally compact and with cocompact automorphism group. Suppose $HDiv(\mathcal{Y}, \mathcal{Z}) \succeq F(r)$ for a pair of non-intersecting hyperplanes \mathcal{Y} and \mathcal{Z} in X . It then follows that $Div(X) \succeq rF(r)$.*

Proof. Let \mathcal{Y}^+ and \mathcal{Z}^+ be half-spaces associated to \mathcal{Y} and \mathcal{Z} such that $\mathcal{Y}^+ \subsetneq \mathcal{Z}^+$.

By the Double Skewering Lemma in [CS11], there exists a $\gamma \in G$ so that $\gamma\mathcal{Z}^+ \subsetneq \mathcal{Y}^+$. Note that $HDiv(\gamma\mathcal{Z}, \mathcal{Z}) \geq F(r)$ since \mathcal{Y} separates \mathcal{Z} from $\gamma\mathcal{Z}$. By Lemma 2.3 in [CS11], γ is hyperbolic and its axis, l , intersects \mathcal{Y} and \mathcal{Z} (γ skewers both \mathcal{Y} and \mathcal{Z}).

Now, we have a chain of equally spaced pairs of hyperplanes $\{\gamma^n\mathcal{Z}, \gamma^{n-1}\mathcal{Z}\}$ along l (isometry moves hyperplanes through l). Hence, the divergence of the geodesic l is at least $rF(r)$. \square

Proof of Theorem 5.3 . The statements in Theorem 5.3 are now an easy consequence of Lemma 5.8, Theorem 5.2 and Theorem 5.10. \square

Remark 5.10.1. We note that Theorem 5.10, Theorem 5.3 and Theorem 6.3 all hold under the different assumption that X is essential, finite-dimensional and $Aut(X)$ has no fixed point at infinity. This is true since the Double Skewering Lemma from [CS11] also holds under these assumptions.

6 Higher Degree Polynomial Divergence in CAT(0) Cube Complexes

We define when a pair of hyperplanes are *degree d k -separated*. We show the divergence of two degree d k -separated hyperplanes is bound below by a polynomial of degree d .

Definition 6.1. Hyperplanes \mathcal{H}_1 and \mathcal{H}_2 are degree 1 k -separated if \mathcal{H}_1 and \mathcal{H}_2 are k -separated.

\mathcal{H}_1 and \mathcal{H}_2 are degree d k -separated if they are k -separated, and for either $i = 1$ or $i = 2$ every geodesic of length k contained in $N(\mathcal{H}_i)$ intersects a pair of degree $(d - 1)$ k -separated hyperplanes.

The main theorem of this section is the following:

Theorem 6.2. *Suppose X is finite-dimensional. If \mathcal{Y} and \mathcal{Z} are degree d k -separated hyperplanes, then $HDiv(\mathcal{Y}, \mathcal{Z})$ is bound below by a polynomial of degree d .*

By combining Theorem 5.10 and 6.2 we immediately get the following:

Theorem 6.3. *Let X be an essential, locally compact CAT(0) cube complex with cocompact automorphism group. If X contains a pair of degree d k -separated hyperplanes, then $Div(X)$ is bound below by a polynomial of degree $d + 1$.*

Proof of Theorem 6.2. The base case, $d = 1$, follows from 2 of Theorem 5.2. For the general case, assume the claim is true for degree $d - 1$ k -separated hyperplanes. Suppose \mathcal{Y} and \mathcal{Z} are degree d k -separated. Let g be a minimal geodesic from \mathcal{Y} to \mathcal{Z} and let $p = g \cap \mathcal{Y}$. Fix $R > 0$, and let α be a path from \mathcal{Y} to \mathcal{Z} which avoids the ball $B_p(R)$. Let D be a combed disk diagram supported by $\{\mathcal{Y}, \alpha, \mathcal{Z}, g^{-1}\}$.

Orient $Y \subset D$ from g to α , and let $A = \{H_1, H_2, \dots, H_n\}$ be dual curves to Y sequentially ordered by the orientation on Y . Since α does not intersect $B_r(p)$, we have that $n \geq r$. Since D is combed and since \mathcal{Y} and \mathcal{Z} are k -separated, it follows for $i > |g| + k$, H_i intersects α . By definition 6.1, there is a subsequence $B = \{K_1, K_2, \dots, K_m\} \subset A$ such that:

1. K_i intersects α
2. For i odd, K_i and K_{i+1} are degree $d - 1$ k -separated.
3. $m \geq \frac{(r-k-|g|)}{k}$
4. $d(K_i, p) \leq ki + |g| + k$

For i odd, let α_i be the segment of α from \mathcal{K}_i to \mathcal{K}_{i+1} . Note that α_i is a path from \mathcal{K}_i to \mathcal{K}_{i+1} which avoids the ball $B_{(r-ki-|g|-k)}(\mathcal{K}_i \cap \mathcal{Y})$. By the induction hypothesis and Lemma 4.4, $|\alpha_i| \geq (r - ki - |g| - k)^{d-1}$. Since we have linearly many segments $\{\alpha_i\}$ whose length is bound below by a degree $d - 1$ polynomial, it follows the length of α is bound below by a degree d polynomial. \square

7 Right-Angled Coxeter Group Divergence

In the next two subsections, we wish to apply the theorems from previous sections to the case of right-angled Coxeter groups (RACGs for short).

Let Γ be the graph associated to a RACG, W_Γ . Let Γ^c be the graph complement of Γ and let I be the set of isolated vertices in Γ^c . i.e., $I = \{v \in V(\Gamma^c) \mid \text{Link}(v) = \emptyset\}$.

I forms a clique in Γ , and Γ is the graph join of the induced subgraph corresponding to I with the induced subgraph corresponding to $\Gamma - I$. Consequently, $W_{(\Gamma-I)}$ is finite index in W_Γ . Divergence is a quasi-isometry invariant, hence divergence results for $W_{(\Gamma-I)}$ apply to W_Γ .

We will from now on assume, without loss of generality, that Γ^c has no isolated vertices for all RACGs considered. The Davis complex for W_Γ under this assumption is essential.

The following definition is a construction used in [DT15].

Definition 7.1 (Γ -complete word). Given a graph Γ which is not a join, let $w_0 = s_1 \dots s_k$ be a word with the property that for every generator $s \in \Gamma$, there exists an i such that $s_i = s$. Furthermore, $m(s_i, s_{i+1}) = \infty$ for all $1 \leq i < k$ and $m(s_1, s_k) = \infty$. Since Γ is not a join, it is always possible to define w_0 , although w_0 is not unique. We call such a word a Γ -complete word and always denote it by w_0 .

We use the definition of a CFS graph used in [BFRHS] and [DT15] (defined below). An *induced square* of a graph Γ is an embedded 4-cycle.

Definition 7.2. Given a graph Γ , define $\square(\Gamma)$ as the graph whose vertices are induced squares of Γ . Two vertices in $\square(\Gamma)$ are adjacent if and only if the corresponding induced

squares in Γ have two non-adjacent vertices in common. For a set of induced squares $S \subset \square(\Gamma)$, define the *support* of S to be all vertices in Γ which are contained in some square in S . Γ is *CFS* if $\square(\Gamma)$ contains a component whose support is $V(\Gamma)$.

Remark 7.2.1. In [BFRHS], the graph join of a CFS graph with a clique graph is still CFS. With the assumption that Γ^c has no isolated vertices, such a graph is not possible and so we omit this from the definition. We note again, however, that the RACG corresponding to a graph that is a join with a clique is commensurable with the RACG corresponding to the graph. So the results in this section still hold in full generality.

7.1 Classification of Linear Divergence

The authors of [BFRHS] classify which right-angled Coxeter groups exhibit linear divergence and the triangle-free case is done in [DT15]. For completeness and as a warm up for the quadratic case, we provide another proof here of this classification.

Theorem 7.3. *If Γ is a join then $Div(W_\Gamma)$ is linear. Otherwise, $Div(W_\Gamma)$ is at least quadratic.*

Proof. Suppose Γ is a join. It follows $W_\Gamma = W_{\Gamma_1} \times W_{\Gamma_2}$. By the assumption that Γ^c has no isolated vertices, both W_{Γ_1} and W_{Γ_2} are infinite. Hence, $Div(W_\Gamma)$ is linear.

Now suppose Γ is not a join. Let $w_0 = s_1 s_2 \dots s_k$ be a Γ -complete word. Let \mathcal{Y} be the hyperplane dual to the letter s_1 in w_0 and \mathcal{Z} the hyperplane dual to the letter s_k in w_0 in the Davis complex of W_Γ . Since, $m(s_1, s_k) = \infty$, it follows \mathcal{Y} and \mathcal{Z} do not intersect. Similarly, any hyperplane dual to the letter s_j in w_0 for $1 < j < k$, does not intersect \mathcal{Y} or \mathcal{Z} .

We will show \mathcal{Y} and \mathcal{Z} are strongly separated. Suppose, for a contradiction, some hyperplane \mathcal{H} intersects both \mathcal{Y} and \mathcal{Z} . Let \mathcal{H} be of type $s \in \Gamma$. It follows that for every j such that $1 \leq j \leq k$, the hyperplane through the letter s_j in w_0 intersects \mathcal{H} . Hence, for every $t \in \Gamma$, $m(t, s) = 2$. But this implies that s is isolated in Γ^c , a contradiction.

Since \mathcal{Y} and \mathcal{Z} are strongly separated, by Theorem 5.3, $Div(W_\Gamma)$ is at least quadratic. \square

7.2 Classification of Quadratic Divergence

We use results from section 5 to classify quadratic divergence in RACGs and show there is a gap between quadratic and cubic divergence in RACGs.

Theorem 7.4. *Suppose Γ is not CFS and is not a join, then W_Γ has divergence greater or equal to a cubic polynomial.*

The proof of the theorem will be given at the end of this subsection. We state the following corollaries which immediately follow.

Corollary 7.4.1. *W_Γ has quadratic divergence if and only if Γ is CFS and is not a join.*

Proof. If Γ is CFS and is not a join, it follows from [BFRHS] that it has quadratic divergence. The other direction follows from Theorem 7.4. \square

Corollary 7.4.2. *If W_Γ is strongly thick of order 2, then W_Γ has cubic divergence.*

Proof. By [BFRHS], W_Γ has at most cubic divergence. Hence, by Theorem 7.4, W_Γ has exactly cubic divergence. \square

The following Lemma guarantees the existence of symbolically k -chain separated hyperplanes when Γ is not CFS.

Lemma 7.5. *Let M be the maximal clique size in Γ . Let $w_0 = s_1 s_2 \dots s_k$ be a Γ -complete word and consider its image in the Davis complex X . Let \mathcal{Y} be the hyperplane dual to w_0 which intersects s_1 and \mathcal{Z} the hyperplane dual to w_0 intersecting s_k . If \mathcal{Y} and \mathcal{Z} are symbolically 2-chain connected then Γ is CFS.*

Proof. Assume \mathcal{Y} and \mathcal{Z} are symbolically 2-chain connected by sequences:

$$\begin{aligned} S_1 &= \{\mathcal{H}^1, \mathcal{K}^1\} \\ S_2 &= \{\mathcal{H}^2, \mathcal{K}^2\} \\ &\dots \\ S_m &= \{\mathcal{H}^m, \mathcal{K}^m\} \end{aligned}$$

Let $a = \{a_1, a_2, \dots, a_m\}$ and $b = \{b_1, b_2, \dots, b_m\}$ be the letters in Γ corresponding respectively to the hyperplanes $\{\mathcal{H}^1, \dots, \mathcal{H}^m\}$ and $\{\mathcal{K}^1, \dots, \mathcal{K}^m\}$. It follows $a \cup b$ forms a CFS subgraph, Δ , of Γ .

Any hyperplane intersecting w_0 cannot intersect \mathcal{Y} or \mathcal{Z} . Thus any such hyperplane separates \mathcal{Y} from \mathcal{Z} . Consequently each hyperplane intersecting w_0 must intersect \mathcal{H}_i and \mathcal{K}_i for some i . Let $L(w_0)$ be the set of generators in the word w_0 , namely $L(w_0) = \{s_1, s_2, \dots, s_k\}$. It follows that for each $s \in L(w_0)$, there exists a j such that s commutes with both $h_j, k_j \in \Delta$.

Given $s \in L(w_0)$, assume $s \notin a \cup b$ as a vertex of Γ , and assume s does not commute with every generator in $a \cup b$. Let $t \in a \cup b$ be such that $m(s, t) = \infty$, $t \in \{a_r, b_r\}$ for some r , and $m(s, a_j) = m(s, b_j) = 2$ for some j with $|r - j| = 1$. This is possible by the above paragraph. It follows $\{s, t, a_j, b_j\}$ forms an induced square which shares two non-adjacent vertices with a square in Δ .

On the other hand, suppose $s_i \in L(w_0)$ commutes with every generator in $A \cup B$. s_{i+1} (if $i = k$ set $s_{i+1} = s_1$) commutes with a_j and b_j for some j . We then have that $\{s_i, s_{i+1}, a_j, b_j\}$ forms an induced square which which shares two non-adjacent vertices with a square in Δ .

We have thus shown every generator in $L(w_0)$ is either contained in Δ or contained in an induced square C which shares two non-adjacent vertices with a square in Δ . Since $L(w_0)$ contains every generator in Γ , we have shown that Γ is CFS. \square

Lemma 7.6. *The Davis complex for W_Γ has 2-alternating geodesics.*

Proof. Choose M to be one larger than the maximal clique size in Γ . Let $g = s_1 s_2 \dots s_M$ be a geodesic of length M with $s_i \in \Gamma$. By Tits' solution to the word problem (see [Dav08]), it follows that $m(s_i, s_j) = \infty$ for some $1 \leq i < j \leq k$. It follows the hyperplane intersecting s_i and the hyperplane intersecting s_j are of non-intersecting type. \square

proof of Theorem 7.4. Theorem 7.4 now follows from the above two lemmas and Theorem 5.3. \square

7.3 Higher Degree Polynomial Divergence in RACGs

In this section, we apply results from section 6 to give graph-theoretic criteria which imply lower bounds on the divergence of a RACG. Together with the machinery of thickness, this allows one to distinguish the exact divergence of many RACGs.

Definition 7.7. Given distinct vertices $s, t \in \Gamma$, (s, t) is a *non-commuting pair* if s is not adjacent to t in Γ .

Definition 7.8. A non-commuting pair (s, t) is rank 1 if s, t are not contained in some induced square of Γ . Additionally (s, t) are rank n if either every non-commuting pair (s_1, s_2) , with $s_1, s_2 \in \text{Link}(s)$, is rank $n - 1$ or every non-commuting pair (t_1, t_2) , with $t_1, t_2 \in \text{Link}(t)$, is rank $n - 1$.




Figure 4: The non-commuting pairs $(4,6)$, $(4,5)$, $(4,9)$, $(5,9)$ and $(6,9)$ for the example graph above are rank 1. It then follows that the non-commuting pair $(7,8)$ is rank 2. Taking this one step further, we see that the non-commuting pair $(9,0)$ is rank 3. By Theorem 7.9, the RACG associated to the above graph has divergence bound below by a polynomial of degree 4. Furthermore, it can easily be checked using techniques from [BHS] that this RACG is thick of order 3 and so the divergence of this group is exactly a quartic polynomial.

Theorem 7.9. *Suppose Γ contains a rank n pair (s, t) , then $\text{Div}(W_\Gamma)$ is bound below by a polynomial of degree $n + 1$.*

Proof. Let M be the maximal clique size in Γ . We claim that hyperplanes of type s and t must be degree n M -separated, in the sense of definition 6.1. If this claim is shown, the theorem follows from Theorem 6.3.

We first prove the base case when $n = 1$. Suppose, for a contradiction, \mathcal{Y} and \mathcal{Z} are of type s and t respectively and that $M + 1$ hyperplanes intersect both \mathcal{Y} and \mathcal{Z} . It follows from Lemma 7.6 that two such hyperplanes, \mathcal{H} and \mathcal{H}' are respectively of type $a, b \in \Gamma$ where (a, b) is a non-commuting pair. However, it then follows $\{s, a, b, t\}$ is an induced square in Γ , contradicting (s, t) being rank 1.

For the general case, suppose (s, t) are rank n and \mathcal{Y} and \mathcal{Z} are hyperplanes of type s and t respectively. Without loss of generality, assume every non-commuting pair (s_1, s_2) , with $s_1, s_2 \in \text{Link}(s)$, are rank $n - 1$. By the induction hypothesis, hyperplanes of type s_1 and type s_2 are degree $n - 1$ M -separated.

Consider any geodesic, $g \subset N(\mathcal{Y})$, of length $M + 1$. By Lemma 7.6, g crosses two hyperplanes of non-commuting type, say of type s_1 and type s_2 . By the above paragraph, (s_1, s_2) must be degree $n - 1$ $M - 1$ -separated. The claim then follows. \square

8 Coxeter Groups

This section explores lower bounds for divergence in Coxeter groups (not necessarily right-angled). Together with the thickness machinery developed for Coxeter groups in [BHS], which provides upper bounds on divergence, we can compute the exact divergence for many of these groups. We use similar arguments to those used in Section 6. We do not make use of a cube complex in this section. Instead, we use the construction of bands in these Van-Kampen diagram which behave similarly to dual curves in CAT(0) cube complex disk diagrams.

In this section, we assume all Coxeter diagrams have at least one edge. Otherwise, W_Γ is virtually trivial or virtually free and exhibits either trivial or infinite divergence.

8.1 Higher Degree Polynomial Divergence

Definition 8.1. Let Γ be a labeled graph. The vertex $v \in \Gamma$ is *r-locally triangle-free* if for all $u \in \Gamma$, such that $d_\Gamma(u, v) < r$, u is not in a triangle. v is *r-locally even* if for all $u \in \Gamma$, such that $d_\Gamma(u, v) < r$, each edge adjacent to u is even labeled or not labeled.

Definition 8.2. For Γ a Coxeter diagram, let L_Γ be the largest integer edge label in Γ . If Γ contains no labeled edges, set $L_\Gamma = 2$.

The following Lemma allows us to choose boundedly spaced generators in a minimal expression for a word $w \in W_{Star(v)}$ such that these generators are not v and do not sequentially coincide.

Lemma 8.3. *Let Γ be a Coxeter diagram. For any $v \in \Gamma$ and any minimal expression, $w = s_1 s_2 \dots s_n$, $s_i \in Star(v)$, for a word $w \in W_{Star(v)}$, there exists a subsequence $\{s_{i_1}, s_{i_2}, \dots, s_{i_m}\}$ such that*

1. $s_{i_j} \neq v$ for all j .
2. $i_1 \leq 2$
3. $i_{j+1} - i_j \leq L_\Gamma$
4. $s_{i_{j+1}} \neq s_{i_j}$ as vertices of Γ
5. $m \geq \frac{n}{L_\Gamma + 1}$.

Proof. Since w is minimal length, there cannot be two v letters appearing consecutively. Hence either the first or second letter is not v . Set t_{i_1} to be this letter. Now note that for any letter $s \in Link(v)$ and $n > L_\Gamma$, we cannot have the expression $susv\dots sv$ or $susv\dots svs$ of length n appearing in w for this would contradict w being reduced. Hence, there is some letter, t_{i_2} not equal to t_{i_1} or v with $i_2 - i_1 \leq L_\Gamma$. We can keep repeating this process, and the lemma follows. \square

Let D be a Van-Kampen diagram for a Coxeter group. Each 2-cell in D has an even number of edges along its boundary path. For a given cell and a given edge along the cell's boundary path, there is a corresponding opposite edge. Furthermore, each edge in D is contained in exactly one cell if it is a boundary edge of D and in exactly two cells if it is not. Two edges e and e' in D are *opposite connected* if there is a sequence of edges $e = e_1, e_2, \dots, e_n = e'$ such that e_i is opposite to e_{i+1} in some 2-cell of D . A

band associated to an edge e in D is the set of all edges opposite connected to e and cells adjacent to these edges.

The construction of bands is utilized in section 1.4 of [Bah05]. There it is also shown that bands do not self-intersect and cannot intersect geodesics twice.

For $u, v \in \Gamma$, an *odd path* from u to v is a path in Γ which only contains edges with odd labels. Let O_v consist of vertices $u \in \Gamma$ for which there is an odd path from u to v . By definition $v \in O_v$.

Let e be an edge in D labeled by some $v \in \Gamma$. It is easy to check the band corresponding to e only contains edges labeled by elements in O_v .

Recall that definition 7.8 of a rank n pair is still valid for Coxeter groups which are not right-angled.

Theorem 8.4. *Let Γ be a Coxeter graph. Suppose (u, v) is a rank n pair. Without loss of generality, we assume that for all distinct $u_1, u_2 \in \text{Link}(u)$, (u_1, u_2) is a rank $n - 1$ pair in Γ . Further assume that u is n -locally triangle free and $n + 1$ -locally even and that v is 1-locally even. It follows that the divergence of the bi-infinite geodesic $\dots uvv\dots$ is bound below by a polynomial of degree $n + 1$.*

The following corollary is immediate:

Corollary 8.4.1. *Let W_Γ be an even Coxeter group such that Γ contains no triangles. If (u, v) is a rank n pair in Γ , then $\text{Div}(W_\Gamma)$ is bound below by a polynomial of degree $n + 1$.*

To prove the above Theorem, we will first need the following technical lemma:

Lemma 8.5. *Let (u, v) be as in Theorem 8.4. Let $g \in W_{\text{Star}(u)}$ and $h \in W_{\text{Star}(v)}$ and $p \in W_\Gamma$ be words written in a minimal length expression. Suppose $|p| \leq L_\Gamma$, $|g| \geq r$ and $|ph| \geq r$. Let α be a shortest path from g to ph in the Cayley graph of W_Γ which does not intersect $B_{\text{id}}(r)$. It follows $|\alpha|$ is bounded below by a polynomial of degree n .*

Proof. The proof will follow by induction on n . We begin with the base case where the rank of (u, v) is $n = 1$. Suppose g is given by the following expression in generators, $g = s_1 s_2 \dots s_l$. Note that $l \geq r$. Let D be a Van-Kampen with boundary path $gah^{-1}p^{-1}$.

Let $T = \{s_{i_1}, s_{i_2}, \dots, s_{i_m}\}$ be a subsequence of $\{s_1, s_2, \dots, s_l\}$ as in Lemma 8.3, and $B = \{B_1, B_2, \dots, B_m\}$ bands in D corresponding to each letter in T . Since u is 2-locally even, each band B_j only contains edges labeled by s_{i_j} . Furthermore, since u is 1-locally triangle free, u is not contained in a triangle. It follows for $i \neq j$, B_i and B_j do not intersect.

At most L_Γ bands can intersect p . Additionally, u and v are rank 1, and so are not in a common square of Γ . It follows for $j > L_\Gamma$, B_j intersects α . Hence, $|\alpha|$ is linear in r , proving the base case.

Now assume the theorem is true for $n - 1$ and that (u, v) are of rank n . The proof proceeds almost the same way as the base case. Consider all the same notation as the base case. For i , such that $L_\Gamma + 1 < i < m$, let α_i be the segment of α between B_i and B_{i+1} , and let p_i be the segment of g from B_i to B_{i+1} . Let g_i be the word along B_i from p_i to α_i , and let h_i be the word along B_{i+1} from p_i to α_i . By the induction hypothesis, $|\alpha_i|$ is bounded below by a polynomial of degree $n - 1$ in $r - i$. Hence, $|\alpha|$ is bounded below by a polynomial of degree n . \square

proof of Theorem 8.4 . Let α be a $B_{id}(r)$ avoidant path from $(uv)^r$ to $(vu)^r$ in the Cayley graph of W_Γ . Let D be a Van-Kampen diagram with boundary path $(uv)^r\alpha(vu)^{-r}$. Since (u, v) is a non-commuting pair in Γ , no pair of bands emanating from the words $(uv)^r$ or from $(vu)^r$ along the boundary path of D can intersect. Hence, each of these bands must intersect α .

Write $(uv)^r$ as $u_1v_1u_2v_2\dots u_rv_r$. Let U_i, V_i be bands corresponding respectively to u_i, v_i . Let D_i be the minimal connected subdiagram of D which includes U_i and V_i . Let α_i be the segment of α contained in D_i . By Lemma 8.5, $|\alpha_i|$ is bound below by a polynomial of degree n in $r - i$. Hence, $|\alpha|$ is bound below by a polynomial of degree $n + 1$. □

8.2 Quadratic Divergence Lower Bound

Definition 8.6. Given an edge labeled Coxeter graph Γ , let $\hat{\Gamma}$ be the graph resulting from homotoping odd labeled edges in Γ to a point. For $v \in \Gamma$ we denote its image in $\hat{\Gamma}$ by $\pi(v)$. Each vertex $\hat{v} \in \hat{\Gamma}$ is labeled by a list, $\pi^{-1}(\hat{v})$. Each edge in $\hat{\Gamma}$ is labeled by the same integer as the corresponding edge in Γ . Note that this new graph can have multiple edges between two vertices.

Theorem 8.7. *Let Γ be a Coxeter graph. If the diameter of $\hat{\Gamma}$ is larger than 2, then W_Γ has at least quadratic divergence.*

Proof. Suppose $d_{\hat{\Gamma}}(\hat{u}, \hat{v}) > 2$ for some $\hat{u}, \hat{v} \in \hat{\Gamma}$. Choose $u \in \pi^{-1}(\hat{u})$ and $v \in \pi^{-1}(\hat{v})$. It follows that $m(u, v) = \infty$. We will show that the bi-infinite geodesic $\dots uvuv\dots$ exhibits quadratic divergence.

Let α be a shortest $2r$ -avoidant path from $(uv)^r$ to $(vu)^r$. Let D be a Van-Kampen diagram with boundary path $(uv)^r\alpha(vu)^{-r}$. Write $(uv)^r = u_1v_1u_2v_2\dots u_rv_r$. Let U_i denote the band emanating from u_i and V_i the band emanating from v_i . Note that for any i, j , U_i cannot intersect V_j . For then, there would be an odd path in Γ from u to some u' , and an odd path from v to some v' , so that $m(u', v') \neq \infty$. However, this would imply $d_{\hat{\Gamma}}(\hat{u}, \hat{v}) \leq 1$, a contradiction.

Fix i . Let D_i denote the minimal connected subdiagram of D containing both U_i and V_i , and let α_i be the subsegment of α contained in D_i . Let $g = s_1\dots s_m$ be the word along the boundary path of U_i from u_i to α_i . It follows $m \geq r - i$. Furthermore, $s_i \in A = \{Star(t) \mid t \in \pi^{-1}(\hat{u})\}$. Note that $d_{\hat{\Gamma}}(\hat{u}, \hat{t}) \leq 1$ for $t \in A$. Let S_j be the band in D_i emanating from s_j . It follows S_j cannot intersect V_i . For then $d_{\hat{\Gamma}}(\hat{u}, \hat{v}) \leq 2$. Hence S_j intersects α_i for each j . It follows, $|\alpha_i|$ is linear in $r - i$. Hence, $|\alpha|$ is quadratic in r . □

Corollary 8.7.1. *Let W_Γ be an even Coxeter group. If $diam(\Gamma) > 2$, then $Div(W_\Gamma)$ is at least quadratic.*

References

- [Ago13] Ian Agol, *The virtual Haken conjecture*, Doc. Math. **18** (2013), 1045–1087. With an appendix by Agol, Daniel Groves, and Jason Manning. ↑2

- [Bah05] Patrick Bahls, *Isomorphism problem in Coxeter groups*, Imperial College Press, 2005. ↑27
- [Beh06] Jason Behrstock, *Asymptotic geometry of the mapping class group and teichmuller space*, *Geometry and Topology* **10** (2006), 1523-1578. ↑1
- [BC12] Jason Behrstock and Ruth Charney, *Divergence and quasimorphisms of right-angled Artin groups*, *Mathematische Annalen* **352** (2012), 339-356. ↑1, 2, 11
- [BFRHS] Jason Behrstock, Victor Falgas-Ravry, Mark F. Hagen, and Timothy Susse, *Global Structural Properties of Random Graphs*. Preprint, arXiv:1505.01913. ↑2, 22, 23, 24
- [BH16] Jason Behrstock and Mark F. Hagen, *Cubulated groups: thickness, relative hyperbolicity, and simplicial boundaries*, *Groups, Geometry, and Dynamics* **10** (2016), no. 2, 649–707. ↑1, 5
- [BHS] Jason Behrstock, Mark F. Hagen, and Alessandro Sisto, *Thickness, relative hyperbolicity, and randomness in Coxeter groups*, with an appendix by Pierre Emmanuel Caprace. Preprint, arXiv:1312.4789. ↑25, 26
- [BD14] Jason Behrstock and Cornelia Druțu, *Divergence, thick groups, and short conjugators*, *Illinois J. Math.* **58** (2014), no. 4, 939–980. ↑1, 3
- [BB05] Anders Bjorner and Francesco Brenti, *Combinatorics of Coxeter groups*, Graduate Texts in Mathematics, vol. 32, Springer, 2005. ↑6
- [BH99] Martin R. Bridson and André Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319, Springer-Verlag, Berlin, 1999. ↑1, 5
- [CS11] Pierre-Emmanuel Caprace and Michah Sageev, *Rank Rigidity for $Cat(0)$ Cube Complexes*, *Geometric and Functional Analysis* **21** (2011), no. 4, 851–891, DOI 10.1007/s00039-011-0126-7. ↑2, 5, 6, 21
- [DJ00] Michael W. Davis and Tadeusz Januszkiewicz, *Right-angled Artin groups are commensurable with right-angled Coxeter groups*, *Journal of Pure and Applied Algebra* **153** (2000), 229-235. ↑2
- [DT15] Pallavi Dani and Anne Thomas, *Divergence in right-angled Coxeter groups*, *Trans. Amer. Math. Soc.* **367** (2015), no. 5, 3549–3577, DOI 10.1090/S0002-9947-2014-06218-1. ↑1, 2, 4, 22, 23
- [Dav08] Michael W. Davis, *The geometry and topology of Coxeter groups*, London Mathematical Society Monographs Series, vol. 32, Princeton University Press, Princeton, NJ, 2008. ↑6, 24
- [DMS10] Cornelia Druțu, Shahar Mozes, and Mark Sapir, *Divergence in lattices in semisimple Lie groups and graphs of groups*, *Trans. Amer. Math. Soc.* **362** (2010), no. 5, 2451–2505, DOI 10.1090/S0002-9947-09-04882-X. ↑5
- [Ger94a] S. M. Gersten, *Quadratic divergence of geodesics in $CAT(0)$ spaces*, *Geom. Funct. Anal.* **4** (1994), 37–51. ↑1
- [Ger94b] ———, *Divergence in 3-manifold groups*, *Geom. Funct. Anal.* **4** (1994), 633–647. ↑2
- [Gro93] Mikhail Gromov, *Geometric Group Theory Volume 2: Asymptotic invariants of infinite groups*, London Mathematical Society Lecture Notes Series, vol. 2, Cambridge University Press, 1993. ↑1
- [GS] Dominik Gruber and Alessandro Sisto, *Infinitely presented graphical small cancellation groups are acylindrially hyperbolic*. Preprint, arXiv:1408.4488. ↑1
- [Gro93] Mikhail Gromov, *Asymptotic invariants of infinite groups*, Geometric group theory, Vol. 2 (Sussex, 1991), 1993, pp. 1–295. ↑1

- [Hag13a] M. F. Hagen, *Weak hyperbolicity of cube complexes and quasi-arboreal groups*, Journal of Topology **7** (2013). ↑7, 8, 9
- [Hag13b] ———, *The simplicial boundary of a $CAT(0)$ cube complex*, Algebraic & Geometric Topology **13** (2013). ↑2
- [KL98] Michael Kapovich and Bernhard Leeb, *3-manifold groups and nonpositive curvature*, GAFA **8** (1998), 841-852. ↑1
- [Mac13] Natasa Macura, *$CAT(0)$ spaces with polynomial divergence of geodesics*, Geometriae Dedicata **163** (2013), 361-378. ↑1
- [OOS09] Alexander Ol'shanskii, Denis Osin, and Mark Sapir, *Lacunary hyperbolic groups*, with an appendix by Michael Kapovich and Bruce Kleiner, Geometry & Topology **13** (2009), 2051-2140. ↑1
- [Wis11] Daniel T. Wise, *The structure of groups with a quasiconvex hierarchy*, 2011. ↑2, 5, 7, 8